arXiv:1409.0691v1 [math.NT] 2 Sep 2014

Genus theory and the factorization of class equations over $\mathbb{F}_p$

Patrick Morton

Aug. 30, 2014

As is well-known, the Hilbert class equation is the polynomial $H_D(X)$ whose roots are the distinct j-invariants of elliptic curves with complex multiplication by the maximal order R_K in the imaginary quadratic field $K = \mathbb{Q}(\sqrt{D})$. A root of $H_D(X)$ generates the Hilbert class field Σ of K over K. The polynomial $H_D(X)$ always has a real root ξ, and over $\mathbb{Q}$ this root generates the real subfield $\Sigma_0 = \mathbb{Q}(\xi)$ of Σ. Recently, Stankewicz [st] found a criterion for the polynomial $H_D(X)$ to have a root (mod p), for a given odd prime p for which the Legendre symbol $(D/p) = -1$. This criterion can be stated as follows.

Theorem (Stankewicz). If p is an odd prime for which $(D/p) = -1$, and p does not divide the discriminant of $H_D(X)$, then $H_D(X)$ has a linear factor over $\mathbb{F}_p$ if and only if

$$\left(\frac{-p}{q}\right) = 1, \quad \forall\ q|D,\ q \text{ an odd prime.}$$

Stankewicz derives this criterion from his analysis of rational p-adic points on twists of Shimura curves. In this note I give a more direct proof of the criterion using genus theory and basic properties of the Hilbert class field. The proof shows that the above theorem fits naturally into a discussion of genus theory. (See [co], [has2], and [ish].)

1 Necessity.

In this and the next section the integer $-N$ will denote the square-free part of the fundamental quadratic discriminant D, and K is the imaginary quadratic field

$K = \mathbb{Q}(\sqrt{-N})$ with discriminant D.

Theorem 1. Let Σ_0 denote the real subfield of the Hilbert class field Σ of the quadratic field $K = \mathbb{Q}(\sqrt{-N})$. Assume p is an odd prime for which $\left(\frac{-N}{p}\right) = -1$. If p has a prime divisor $\mathfrak{p}$ of degree 1 in Σ_0, then

$$\left(\frac{-p}{q}\right) = 1, \quad \forall\ q|N,\ q \text{ an odd prime.}$$

To prove this we use the decomposition

$$D = \prod_{q|D} q^*,\ q^* = (-1)^{(q-1)/2}q,\ q \text{ odd},\ 2^* = -4, 8, -8,$$

where the product is over all the prime divisors of D.

The genus field of K is the field Ω, which is obtained by adjoining all the square-roots $\sqrt{q^*}$ to K, as q varies over the prime divisors of D. It is the largest unramified extension of K which is abelian over $\mathbb{Q}$, so that $\Omega \subseteq \Sigma$.

Assume that the odd prime p has a first degree prime divisor in Σ_0. The conditions $\left(\frac{-N}{p}\right) = -1$ and p odd imply that p does not divide D, and p has a first degree prime divisor in every subfield of Σ_0. If q is a prime $\equiv 1 \pmod 4$, then $q^* = q$, so $\mathbb{Q}(\sqrt{q}) \subseteq \Sigma_0$. Hence, p has a first degree prime divisor in $\mathbb{Q}(\sqrt{q})$, which implies that

$$\left(\frac{q}{p}\right) = 1,\ q \equiv 1 \pmod 4,\ q|N. \tag{1.1}$$

This implies then that

$$\left(\frac{-p}{q}\right) = 1,\ q \equiv 1 \pmod 4,\ q|N. \tag{1.2}$$

If $2|D$ and $2^* = 8$, the same argument also gives $\left(\frac{2}{p}\right) = 1$, so $p \equiv \pm 1 \pmod 8$.

On the other hand, if there are several primes $q_i \equiv 3 \pmod 4$, $i = 1, 2$, then $\sqrt{q_i^*} \in \Sigma$ implies that $\sqrt{q_1^* q_2^*} = \sqrt{q_1 q_2} \in \Sigma_0$. Then p has a first degree prime divisor in $\mathbb{Q}(\sqrt{q_1 q_2})$, so we have

$$\left(\frac{q_1 q_2}{p}\right) = 1,\ q_i \equiv 3 \pmod 4,\ q_i|N.$$

It follows that

$$\left(\frac{q_1^*}{p}\right) = \left(\frac{q_2^*}{p}\right), \ q_1 \equiv q_2 \equiv 3 \pmod 4, \ q_i | N. \tag{1.3}$$

We get a similar conclusion when $2|D$ and $2^* = -4, -8$, namely

$$\left(\frac{2^*}{p}\right) = \left(\frac{q^*}{p}\right), \ 2^* = -4, -8, \ q \equiv 3 \pmod 4, \ q|N. \tag{1.4}$$

Now we use the fact that

$$\left(\frac{D}{p}\right) = \left(\frac{-N}{p}\right) = \prod_{q|D} \left(\frac{q^*}{p}\right) = -1. \tag{1.5}$$

From (1.1), (1.3), (1.4) the terms with $q \equiv 1 \pmod 4$ or $q^* = 8$ drop out, and we are left with

$$\left(\frac{q^*}{p}\right)^r = -1,$$

where r is the number of prime divisors of D with $q \equiv 3 \pmod 4$ or $q^* = -4, -8$. But this implies that r is odd and $\left(\frac{q^*}{p}\right) = -1$ for all these prime divisors. Hence,

$$\left(\frac{-p}{q}\right) = 1, \text{ if } q \equiv 3 \pmod 4, \ q|N.$$

Together with (1.2), this proves Theorem 1. □

Corollary 1. If p does not divide the discriminant of $H_D(X)$, $\left(\frac{-N}{p}\right) = -1$, and $H_D(X) \pmod p$ has a root in $\mathbb{F}_p$, then

$$\left(\frac{-p}{q}\right) = 1, \quad \forall \ q|N, \ q \text{ an odd prime.}$$

Proof. Since p does not divide the discriminant of $H_D(X)$ and a real root of $H_D(X)$ generates Σ_0, it is clear that the factors of $H_D(X) \bmod p$ correspond 1-1 to the prime divisors of p in Σ_0. The corollary is now immediate from Theorem 1. □

2 Sufficiency.

Now we prove the converse of Theorem 1:

Theorem 2. Let Σ_0 denote the real subfield of the Hilbert class field Σ of the quadratic field $K = \mathbb{Q}(\sqrt{-N})$. Assume p is an odd prime for which $\left(\frac{-N}{p}\right) = -1$. If p satisfies the condition

$$\left(\frac{-p}{q}\right) = 1, \quad \forall\, q|N,\ q \text{ an odd prime},$$

then p has a prime divisor $\mathfrak{p}$ of degree 1 in Σ_0. $\square$

To prove this we consider the decomposition group of a prime divisor $\mathfrak{P}$ of p in Σ. First we note that if $\left(\frac{-p}{q}\right) = 1$ for all odd prime divisors q of N, then (1.1) holds, as does

$$\left(\frac{q^*}{p}\right) = -1,\ q \equiv 3 \pmod 4,\ q|N,\ q \text{ prime}. \tag{2.1}$$

Now (1.5) implies that

$$\left(\frac{2^*}{p}\right) = (-1)^{r-1} \cdot (-1) = (-1)^r, \text{ if } 2|D,$$

where $r-1$ is the number of primes $q \equiv 3 \pmod 4$ dividing N. But if $2|D$, then either:

$N \equiv 1 \pmod 4$, in which case $2^* = -4$ and $r-1$ is even, so that r is odd, implying that $\left(\frac{2^*}{p}\right) = \left(\frac{-4}{p}\right) = -1$;

$N \equiv 2 \pmod 8$, in which case $2^* = -8$ and r is again odd, giving $\left(\frac{2^*}{p}\right) = \left(\frac{-8}{p}\right) = -1$;

or $N \equiv 6 \pmod 8$, in which case $2^* = 8$ and $r-1$ is odd, giving $\left(\frac{2^*}{p}\right) = \left(\frac{8}{p}\right) = 1$.

Thus, if $2|D$, we have (1.4) and the assertion in the sentence following (1.2). This shows that p splits completely in the real subfield Ω_0 of the genus field Ω. (Note

that $[\Omega : K] = 2^{t-1}$, where t is the number of distinct prime factors of D. Thus $[\Omega_0 : \mathbb{Q}] = 2^{t-1}$, as well.) Hence, the decomposition field of any prime divisor $\mathfrak{P}$ of p in Σ contains the field Ω_0, and therefore the decomposition group $G_{\mathfrak{P}}$ is contained in $H = Gal(\Sigma/\Omega_0)$.

I claim that it suffices to show $G_{\mathfrak{P}} = \{1, \tau\}$ for some $\mathfrak{P}|p$, where τ is complex conjugation. If this holds, then Σ_0, which is the fixed field of τ, is the largest field in which the prime below $\mathfrak{P}$ has degree 1, i.e. $\mathfrak{p} = \mathfrak{P}\mathfrak{P}^{\tau}$ is a first degree prime divisor of p in Σ_0.

Let $J = H \cap Gal(\Sigma/K)$ be the subgroup of $Gal(\Sigma/K)$ corresponding to Ω in the Galois correspondence, so that $[H : J] = 2$ and $H = J \cup J\tau$. By the genus theory [h], J corresponds to the subgroup of squares in $Pic(R_K)$, in the Artin correpondence between ideal classes in R_K and elements of the Galois group $Gal(\Sigma/K)$.

We now have what we need to complete the proof. The decomposition group $G_{\mathfrak{P}}$ is a subgroup of H of order 2. This is because p is inert in K, so that $(p) = pR_K$ is a principal ideal and therefore p splits completely in the extension Σ/K. Furthermore, we know that K is not contained in the decomposition field of any $\mathfrak{P}$, and therefore $G_{\mathfrak{P}} \not\subset J \subset Gal(\Sigma/K)$. Hence, $G_{\mathfrak{P}} \subset H$ is generated by some $\sigma\tau$, with $\sigma \in J$. But $\sigma = \psi^2$ for some $\psi \in Gal(\Sigma/K)$, by the characterization of the group J, and $\psi^{-1}G_{\mathfrak{P}}\psi = G_{\mathfrak{P}^{\psi}} = \{1, \psi^{-1}\sigma\tau\psi\}$, with $\psi^{-1}\sigma\tau\psi = \psi^{-2}\sigma\tau = \tau$. This shows that $G_{\mathfrak{P}^{\psi}} = \{1, \tau\}$ and completes the proof. $\square$

Corollary 2. If p satisfies $\left(\frac{-N}{p}\right) = -1$ and $\left(\frac{-p}{q}\right) = 1$, for all odd primes q such that $q|N$, then $H_D(X) \pmod p$ has a root in $\mathbb{F}_p$.

Proof. Theorem 2 implies that $H_D(X)$ has a linear factor over $\mathbb{Q}_p$ and therefore $H_D(X) \pmod p$ has a root in $\mathbb{F}_p$. $\square$

The proof of Theorem 2 shows that when p has a first degree prime divisor $\mathfrak{p}$ in Σ_0, then it has as many prime divisors of degree 1 as there are distinct elements ψ in $Gal(\Sigma/K)$ for which $G_{\mathfrak{P}^{\psi}} = \psi^{-1}G_{\mathfrak{P}}\psi = \{1, \tau\}$, where $\mathfrak{P}|\mathfrak{p}$. This holds if and only if $\psi^{-1}\tau\psi = \tau$, i.e., if and only if $\psi^2 = 1$. The number of such elements ψ is exactly 2^{t-1}, since this is the order of the 2-Sylow subgroup of the class group $Pic(R_K)$. Thus we have:

Theorem 3. If p is an odd prime for which $\left(\frac{-N}{p}\right) = -1$, and p has a prime divisor of degree 1 in Σ_0, then it has exactly 2^{t-1} such prime divisors, where t is the number of distinct prime factors of D. $\square$

Taken together, Theorems 1-3 yield the following Decomposition Law for the real subfield Σ_0 of Σ.

Prime Decomposition Law in Σ_0. Let p be an odd prime that does not divide D.

(a) If $\left(\frac{D}{p}\right) = 1$, then in Σ_0, p splits into $h(D)/f$ primes of degree f over $\mathbb{Q}$, where f is the order of a prime ideal divisor $\wp$ of p in $Pic(R_K)$.

(b) If $\left(\frac{D}{p}\right) = -1$ and $\left(\frac{-p}{q}\right) = 1$ for all odd prime divisors q of D, then in Σ_0, p splits into $r_1 = 2^{t-1}$ primes of degree 1 and $r_2 = (h(D) - 2^{t-1})/2$ primes of degree 2 over $\mathbb{Q}$.

(c) If $\left(\frac{D}{p}\right) = -1$ and $\left(\frac{-p}{q}\right) = -1$ for some odd prime divisor q of D, then in Σ_0, p splits into $h(D)/2$ primes of degree 2 over $\mathbb{Q}$. $\square$

This law immediately implies the following density result.

Theorem 4. The density of primes $p \in \mathbb{Z}^+$ for which $H_D(X)$ has a linear factor (mod p) is $d(P(\Sigma_0)) = \frac{1}{2h(D)} + \frac{1}{2^t}$, where t is the number of distinct prime factors of D.

References.

[co] D. A. Cox, *Primes of the form $x^2 + ny^2$*, John Wiley and Sons, New York, 1989.

[hak1] F. Halter-Koch, Arithmetische Theorie der Normalkörper von 2-Potenzgrad mit Diedergruppe, J. Number Theory 3 (1971), 412-443.

[hak2] F. Halter-Koch, Geschlechtertheorie der Ringklassenkörper, J. reine angew. Math. 250 (1971), 107-108.

[hak3] F. Halter-Koch, Eine allgemeine Geschlechtertheorie und ihre Anwendung auf Teilbarkeitsaussagen für Klassenzahlen algebraischer Zahlkörper, Math. Annalen 233 (1978), 55-63.

[has1] H. Hasse, *Bericht über neuere Untersuchungen und Probleme aus der Theorie der algebraischen Zahlkörper*, I, Ia, II, Physica Verlag, Würzburg-Vienna, 1970.

[has2] H. Hasse, Zur Geschlechtertheorie in quadratischen Zahlkörpern, J. Math. Soc. Japan 3 (1951), 45-51.

[ish] M. Ishida, The Genus Fields of Algebraic Number Fields, Lecture Notes in Mathematics 555, Springer, Berlin, 1976.

[st] J. Stankewicz, Twists of Shimura curves, Canad. J. Math. 66 (2014), 924-960; and at http://arxiv.org/abs/1208.3594.

Dept. of Mathematical Sciences

Indiana University - Purdue University at Indianapolis (IUPUI)

402 N. Blackford St., Indianapolis, IN, USA 46202

e-mail: pmorton@math.iupui.edu